

No part of the candidate's evidence in this exemplar material may be presented in an external assessment for the purpose of gaining an NZQA qualification or award.

SUPERVISOR'S USE ONLY

S

93202



Draw a cross through the box (☒) if you have NOT written in this booklet

☐

+

OUTSTANDING SCHOLARSHIP EXEMPLAR



Mana Tohu Mātauranga o Aotearoa
New Zealand Qualifications Authority

Scholarship 2024 Calculus

Time allowed: Three hours
Total score: 32

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

You should answer ALL the questions in this booklet.

Pull out Formulae and Tables Booklet S–CALCF from the centre of this booklet.

Show ALL working. Correct answers only will not be sufficient.


Check that this booklet has pages 2–24 in the correct order and that none of these pages is blank.

Do not write in any cross-hatched area (▨). This area will be cut off when the booklet is marked.

YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.

QUESTION ONE

- (a) Consider the curve $y = \frac{2x^2 + 1}{3x^2 - 4x - 2}$, which is shown below.



- (i) Find the coordinates of any stationary points on the curve, and determine their nature.

$$y = \frac{2x^2 + 1}{3x^2 - 4x - 2} \quad \text{stationary points.}$$

$$x = \frac{1}{4}, y = -\frac{2}{5}$$

$$\frac{dy}{dx} = \frac{4x(3x^2 - 4x - 2) - (2x^2 + 1)(6x - 4)}{(3x^2 - 4x - 2)^2}$$

$$x = -2, y = \frac{1}{2}$$

$$= \frac{12x^3 - 16x^2 - 8x - 12x^3 + 8x^2 - 6x + 4}{(3x^2 - 4x - 2)^2}$$

$$= \frac{-8x^2 - 14x + 4}{(3x^2 - 4x - 2)^2} = 0$$

$$4x^2 + 7x - 2 = 0$$

$$(4x - 1)(x + 2) = 0$$

$$x = \frac{1}{4}, -2$$

$$\frac{d^2y}{dx^2} = \frac{(-16x - 14)(3x^2 - 4x - 2)^2 - (-8x^2 - 14x + 4)2(3x^2 - 4x - 2)(6x - 4)}{(3x^2 - 4x - 2)^4}$$

- (ii) Find the value of $\lim_{x \rightarrow \infty} \frac{2x^2 + 1}{3x^2 - 4x - 2}$.

Hence, find the coordinate(s) where the curve intersects its own asymptote(s).

$$\lim_{x \rightarrow \infty} \frac{2x^2 + 1}{3x^2 - 4x - 2} = \lim_{x \rightarrow \infty} \frac{2x^2 - \frac{8}{3}x - \frac{4}{3} + \frac{8}{3}x + \frac{7}{3}}{3x^2 - 4x - 2}$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{2}{3}(3x^2 - 4x - 2) + \frac{8}{3}x + \frac{7}{3}}{3x^2 - 4x - 2}$$

$$= \lim_{x \rightarrow \infty} \left(\frac{2}{3} + \frac{\frac{8}{3}x + \frac{7}{3}}{3x^2 - 4x - 2} \right) = \frac{2}{3}$$

$\therefore$ Curve has a horizontal asymptote $y = \frac{2}{3}$

When curve intersects asymptote,

$$\frac{2}{3} = \frac{2x^2 + 1}{3x^2 - 4x - 2}$$

$$6x^2 - 8x - 4 = 6x^2 + 3$$

$$8x = -7$$

$$x = -\frac{7}{8}$$

$$y = \frac{2}{3}$$

$\therefore$

Curve intersects asymptote at $(-\frac{7}{8}, \frac{2}{3})$

- (b) For what real value(s) of α does the following system of equations have no real solutions?

$$x^3 + y^3 = 2$$

$$x + y = \alpha$$

$$y = \alpha - x$$

$$x^3 + (\alpha - x)^3 = 2$$

$$x^3 + \alpha^3 - 3\alpha^2x + 3\alpha x^2 - x^3 = 2$$

$$3\alpha x^2 - 3\alpha^2x + \alpha^3 - 2 = 0$$

Taking the discriminant,

$$(-3\alpha^2)^2 - 4(3\alpha)(\alpha^3 - 2) < 0$$

$$12\alpha^4 - 12\alpha^4 + 24\alpha < 0$$

$$-12\alpha^4 + 24\alpha < 0$$

$$4\alpha(3 - \alpha^3) < 0$$

$$3\alpha(4 - \alpha^3) > 0$$

Taking discriminant:

$$(-3\alpha^2)^2 - 4(3\alpha)(\alpha^3 - 2) < 0$$

$$9\alpha^4 - 12\alpha^4 + 24\alpha < 0$$

$$-3\alpha^4 + 24\alpha < 0$$

$$3\alpha(\alpha^3 - 8) > 0$$

$$\alpha > 2, \alpha < 0$$

- (c) A landscaper is building steps up one side of a 48 metre high hill.

The slope of this side of the hill can be modelled by the curve


$$y = kx^2(126 - x) = 126kx^2 - kx^3$$

where both x and y are in metres and k is a constant to be determined.

Building regulations state that each step needs to have a **minimum** depth of 280 mm and a **maximum** height of 190 mm, as shown.

If the landscaper chooses to build each step with a depth of 280 mm, would the steps all satisfy the maximum height regulations?

You must use calculus to support your answer.



~~W~~ Hill has a maximum at 48 meters height

$$\frac{dy}{dx} = 2kx(126 - x) - kx^2$$

$$-kx^2 + 252kx = 0 \Rightarrow -3kx^2 + 252kx = 0$$

$$kx(252 - 3x) = 0$$

$$x = \frac{252}{3}, 0$$

$$x = 0 \text{ is foot of hill, } \therefore x = \frac{252}{3}$$

$$x = \frac{252}{3}, y = 48, 48 = k \left(\frac{252}{3} \right)^2 (126 - \frac{252}{3}) \Rightarrow k = \frac{48}{\left(\frac{252}{3} \right)^2 (126 - \frac{252}{3})} = \frac{1}{6174}$$

The height of - Let h_n be height of the n th step

$$h_n = \frac{1}{6174} ((0.28n)^2 (126 - (0.28n)) - (0.28(n-1))^2 (126 - (0.28(n-1))))$$

Substituting $n=150$,

$$\begin{aligned} h_n &= \frac{1}{6174} ((0.28 \times 150)^2 (126 - 0.28 \times 150) - (0.28 \times 149)^2 (126 - 0.28 \times 149)) \\ &= 0.23999 \dots \text{ m} \\ &= 239.99 \dots \text{ mm} > 190 \end{aligned}$$

$\therefore$ Not all steps satisfy maximum height regulations

QUESTION TWO


- (a) For any two functions f and g defined on an interval, $a \leq x \leq b$, the **max value** function is defined as:

$$\max(f, g) = \begin{cases} f(x) & \text{if } f(x) \geq g(x) \\ g(x) & \text{if } f(x) < g(x) \end{cases}$$

To demonstrate, an example is shown opposite.

The max value function can be applied to more than two functions in a similar manner.

Consider the function $h(x) = \max(2\sqrt{x}, 2x, x^2)$ on the interval $0 \leq x \leq 3$.



Evaluate the definite integral $\int_0^3 h(x) dx$.

When $2\sqrt{x} = 2x$, $\sqrt{x} = x$, $x^2 = x$, $x^2 - x = 0$, $x = 0, 1$
 For $0 \leq x \leq 1$, $2\sqrt{x} > 2x$
 when $2\sqrt{x} > 2x$, $\sqrt{x} > x$, $x > x^2$, $x^2 - x < 0$, $0 < x < 1$
 when $2\sqrt{x} > x^2$ if $2x > x^2$, $x^2 - 2x < 0$, $0 < x < 2$

$$\therefore h(x) = \begin{cases} 2\sqrt{x} & \text{for } 0 \leq x \leq 1 \\ 2x & \text{for } 1 \leq x < 2 \\ x^2 & \text{for } 2 \leq x \leq 3 \end{cases}$$

$$\begin{aligned} \therefore \int_0^3 h(x) dx &= \int_0^1 2x^{\frac{1}{2}} dx + \int_1^2 2x dx + \int_2^3 x^2 dx \\ &= \left[\frac{1}{3} x^{\frac{3}{2}} \right]_0^1 + \left[x^2 \right]_1^2 + \left[\frac{1}{3} x^3 \right]_2^3 \\ &= \frac{1}{3} \times 1^{\frac{3}{2}} - \frac{1}{3} \times 0^{\frac{3}{2}} + 2^2 - 1^2 + \frac{1}{3} \times 3^3 - \frac{1}{3} \times 2^3 \\ &= \frac{29}{3} \end{aligned}$$

- (b) Given that $\frac{dy}{dx} = 1 + y^2$, find the value of M for which $\frac{d^3y}{dx^3} = M(1 + y^2)(1 + 3y^2)$.

$$\frac{dy}{dx} = 1 + y^2$$

$$\frac{d^2y}{dx^2} = 2y \frac{dy}{dx}$$

$$\frac{d^3y}{dx^3} = 2 \frac{dy}{dx} + 2y \frac{d^2y}{dx^2}$$

$$= 2 \frac{dy}{dx} + 2y \times 2y \frac{dy}{dx}$$

$$= 2(1 + 2y^2) \frac{dy}{dx}$$

$$= 2(1 + 2y^2)(1 + y^2)$$

$$M = 2$$

- (c) For any real number x , the **absolute value** of x is defined as:

$$|x| = \begin{cases} -x & \text{if } x < 0 \\ x & \text{if } x \geq 0 \end{cases}$$

Consider the function $s_n(x) = |x-1| + |x-2| + \dots + |x-n|$ with $n > 2$.

- (i) Find $s'_3\left(\frac{5}{2}\right)$, the derivative of $s_3(x)$ evaluated at $x = \frac{5}{2}$.
 $\text{at } x = \frac{5}{2}, x-2 \geq 0, x-3 < 0$

$$s_3(x) = |x-1| + |x-2| + |x-3|$$

$$= x-1 + x-2 - x+3 = x$$

$$s'_3(x) = 1$$

$$s'_3\left(\frac{5}{2}\right) = 1$$

- (ii) Find the minimum value of $s_{2024}(x)$.

Justify your answer.

$$s_2(x) = |x-1| + |x-2|$$

$$x < 1: s_2 = -x+1 -x+2 = -2x+3$$

$$1 < x < 2: s_2(x) = -x+1 + x-2 = -1$$

$$x > 2: s_2(x) = x-1 + x-2 = 2x-3$$

For $s_{2024}(x)$, $s_n(x)$ must be defined by $n+1$ different sections. Let $f_n(x) = |x-n|$, $f'_n(x) = \pm 1$

$$s'_{2024}(x) = \sum_{n=1}^{2024} f'_n(x)$$

As x increases, $s'_{2024}(x)$ increases as each $f'_n(x)$ becomes equal to 1

$s_{2024}(x)$ is continuous

$\therefore$ Minimum occurs when

Let $1012 \leq x \leq 1013$, i.e. $x = 1012 + \alpha$, where $0 \leq \alpha \leq 1$

$$s_{2024}(x) = 1012 + \alpha - 1 + 1012 + \alpha - 2 + 1012 + \alpha - 3 + \dots + 1012 + \alpha - 1012 + 1013 - 1012 - \alpha + 1014 - 1012 - \alpha + \dots + 2024 - 1012 - \alpha$$

$$= 1013 + 1014 + \dots + 2024 - 1 - 2 - \dots - 1012 = 1012^2$$

QUESTION THREE

- (a) Find the exact value of $\int_0^{\frac{\pi}{4}} \frac{\cos 2x}{\cos^2 x} dx$.

$$\int_0^{\frac{\pi}{4}} \frac{\cos 2x}{\cos^2 x} dx = \int_0^{\frac{\pi}{4}} \frac{2\cos^2 x - 1}{\cos^2 x} dx$$

$$= \int_0^{\frac{\pi}{4}} \left(2 - \frac{1}{\cos^2 x} \right) dx$$

$$= \int_0^{\frac{\pi}{4}} (2 - \sec^2 x) dx$$

$$= \left[2x - \tan x \right]_0^{\frac{\pi}{4}}$$

$$= 2 \times \frac{\pi}{4} - \tan \frac{\pi}{4} - 2 \times 0 + \tan 0$$

$$= \frac{\pi}{2} - 1$$


- (b) Alice takes a ride at an amusement park that involves
- five spinning teacups that rotate on circular saucers
 - a circular platform that rotates around a large teapot at its centre.

The coordinates of Alice's motion in the xy -plane (shown as point A in the diagram below) can be described by the parametric equations:

$$x = 4\cos\theta + \cos 4\theta$$

$$y = 4\sin\theta + \sin 4\theta$$

where θ is the anti-clockwise angle shown.



<https://www.flickr.com/photos/nycedc/4691066745>

- (i) Find the coordinates in the first quadrant where Alice is closest to the teapot.

Justify your answer.

~~dist~~ let dist = r

$$r = \sqrt{x^2 + y^2}$$

$$= \sqrt{16\cos^2\theta + 8\cos\theta\cos 4\theta + \cos^2 4\theta + 16\sin^2\theta + 8\sin\theta\sin 4\theta + \sin^2 4\theta}$$

$$= \sqrt{16 + 8\cos\theta\cos 4\theta + 8\sin\theta\sin 4\theta}$$

$$r^2 = 17 + 8\cos\theta\cos 4\theta + 8\sin\theta\sin 4\theta$$

$$\frac{dr^2}{d\theta} = 8\cos\theta - 8\sin\theta\cos 4\theta - 32\cos\theta\sin 4\theta + 8\cos\theta\sin 4\theta + 32\sin\theta\cos 4\theta$$

$$= 24\sin\theta\cos 4\theta - 24\cos\theta\sin 4\theta = 0$$

$$2\sin\theta\cos 4\theta - 2\cos\theta\sin 4\theta = 0$$

$$\sin 5\theta + \sin(-3\theta) - \sin 5\theta + \sin(-3\theta) = 0$$

$$\sin(-3\theta) = 0 \quad \theta = 0, \frac{\pi}{3}$$

- (ii) Find the distance Alice travels in one complete rotation around the teapot.

Note: when a curve is defined parametrically by the equations $x = f(\theta)$ and $y = g(\theta)$ on an interval $\alpha \leq \theta \leq \beta$, we can find its length, L , by using the formula:

$$L = \int_{\alpha}^{\beta} \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta$$

$$\frac{dx}{d\theta} = -4\sin\theta - 4\sin 4\theta$$

$$\frac{dy}{d\theta} = 4\cos\theta + 4\cos 4\theta$$

$$L = \int_0^{2\pi} \sqrt{(-4\sin\theta - 4\sin 4\theta)^2 + (4\cos\theta + 4\cos 4\theta)^2} d\theta$$

$$= \int_0^{2\pi} \left(16\sin^2\theta + 32\sin\theta\sin 4\theta + 16\sin^2 4\theta + 16\cos^2\theta + 32\cos\theta\cos 4\theta + 16\cos^2 4\theta \right)^{\frac{1}{2}} d\theta$$

$$= 32 \int_0^{2\pi} (1 + 1 + 32\sin\theta\sin 4\theta + 32\cos\theta\cos 4\theta)^{\frac{1}{2}} d\theta$$

$$= 32 \int_0^{2\pi} (2 + 32\sin\theta\sin 4\theta + 32\cos\theta\cos 4\theta)^{\frac{1}{2}} d\theta$$

$$= 4 \int_0^{2\pi} (2 + 2\sin\theta\sin 4\theta + 2\cos\theta\cos 4\theta)^{\frac{1}{2}} d\theta$$

$$= 4 \int_0^{2\pi} (2 + \cos(-3\theta) - \cos(5\theta) + \cos(5\theta) + \cos(-3\theta))^{\frac{1}{2}} d\theta$$

$$= 4 \int_0^{2\pi} (2 + 2\cos(3\theta))^{\frac{1}{2}} d\theta$$

$$= 4 \int_0^{2\pi} (2 + 2\cos^2 \frac{3}{2}\theta - 2\sin^2 \frac{3}{2}\theta)^{\frac{1}{2}} d\theta$$


$$= 4 \int_0^{2\pi} (2\sin^2 \frac{3}{2}\theta + 2\cos^2 \frac{3}{2}\theta + 2\cos^2 \frac{3}{2}\theta - 2\sin^2 \frac{3}{2}\theta)^{\frac{1}{2}} d\theta$$

$$= 4 \int_0^{2\pi} 2\cos \frac{3}{2}\theta d\theta = 4 \left[\frac{4}{3} \sin \frac{3}{2}\theta \right]_0^{2\pi} =$$

- (c) In 2023, two American high-school students discovered a new proof of the Pythagorean Theorem using trigonometry. In this question we will work through the key steps to derive their result.

The students' proof makes use of the diagram below, which consists of an infinite number of **similar** right-angled triangles enclosed within a large right-angled triangle. It was referred to by the students as a "waffle cone".

Note: to avoid circular logic, you should **not** make use of the Pythagorean Theorem or any of the Pythagorean trigonometric identities at any step in your working for this question.



- (i) Show that $c^2 = \frac{2ab}{\sin 2\alpha}$.

$$\text{Area of } ABC = \frac{1}{2} b \times 2a = \frac{1}{2} c \times c \times \sin 2\alpha$$

$$ab = \frac{1}{2} c^2 \sin 2\alpha$$

$$c^2 = \frac{2ab}{\sin 2\alpha}$$

Triangle formula is independent of pythagorean theorem

- (ii) Show that $u = \frac{2abc}{b^2 - a^2}$ and find a similar expression for v .

Hence, prove that $a^2 + b^2 = c^2$.

Hint: the following formulae will prove useful:

$$T_n = T_1 r^{n-1} \quad S_n = T_1 \left(\frac{1-r^n}{1-r} \right) \quad \text{along } u$$

Let the ~~first~~ ^{nth} hypotenuse be T_n

$$T_1 = \sqrt{(2a)^2 + (2b)^2} = 2\sqrt{a^2 + b^2} = 2c \quad \text{quadrilateral}$$

~~suppose~~ we have a ~~parallelogram~~ ^{quadrilateral} $WXZY$
~~similar to~~ $\triangle CED$, where $WX = x$

$$XY = x \times \frac{a}{b}$$

$$ZY = XY \times \frac{a}{b} = x \times \left(\frac{a}{b}\right)^2$$

Each subsequent quadrilateral in the cone has their side lengths multiplied by $\left(\frac{a}{b}\right)^2$

$$\therefore T_n = \frac{2ac}{b} \times \left(\frac{a^2}{b^2}\right)^{n-1}$$

$$u = S_\infty = \frac{2ac}{b} \left(\frac{1 - \left(\frac{a^2}{b^2}\right)^\infty}{1 - \frac{a^2}{b^2}} \right) \quad \text{2abc}$$

$$= \frac{\frac{2ac}{b}}{1 - \frac{a^2}{b^2}} = \frac{2abc}{b^2 - a^2}$$

The first hypotenuse along $v = c$, ~~let the nth be~~ U_n

$$\therefore U_1 = c, U_n = c \left(\frac{a^2}{b^2}\right)^{n-1} \quad \text{but is not part of a}$$

$$v = S_\infty = c \left(\frac{1 - \left(\frac{a^2}{b^2}\right)^\infty}{1 - \frac{a^2}{b^2}} \right) = \frac{cb^2}{b^2 - a^2} \quad \text{similar quadrilateral}$$

$$\sin 2\alpha = \frac{u}{v} = \frac{\frac{2abc}{b^2 - a^2}}{\frac{cb^2}{b^2 - a^2}} = \frac{2a}{b} = \frac{2ab}{c^2}$$

$$\text{Second} = c \div b \times \frac{2a^2}{b} = \frac{2ca^2}{b^2}$$

$$\therefore v = c + S_\infty = c + \frac{2ca^2}{b^2 - a^2} = c + \frac{2ca^2}{b^2 - a^2}$$


QUESTION FOUR

(a) Consider all the complex numbers z that satisfy all of the following three conditions:

- $-\frac{\pi}{3} \leq \arg(z) \leq \frac{\pi}{3}$
- $z + \bar{z} \leq 4$ $2x \leq 4$
- $|z| \geq 2$

Find the exact area of the region generated in an Argand diagram by the locus of points that represent z .

Use the Argand diagram below to support your working.



$$\text{Area} = \text{Triangle} - \text{sector}$$

$$= \frac{1}{2} \times 2 \times 4\sqrt{3} - \frac{2\pi}{3} \times 2^2 \times \frac{1}{2}$$

$$= 4\sqrt{3} - \frac{4\pi}{3}$$

(b) If $z = \cos\theta + i\sin\theta$ and $0 \leq \theta \leq \frac{\pi}{2}$, prove the following:

(i) $\frac{1}{1-z\cos\theta} = 1 + i\cot\theta$

$$1 + \frac{\cos\theta}{\sin\theta}i = \frac{\sin\theta + \cos\theta i}{\sin\theta}$$

$$\frac{1}{1-z\cos\theta} = \frac{1}{1-\cos^2\theta - \cos\theta i \sin\theta} = \frac{1}{1-\cos^2\theta - \sin\theta\cos\theta i}$$

$$= \frac{1}{\sin^2\theta - \sin\theta\cos\theta i}$$

$$1 + i\cot\theta = 1 + \frac{i\cos\theta}{\sin\theta} = 1 - \frac{\cos\theta}{i\sin\theta}$$

$$\frac{1}{1-z\cos\theta} = \frac{1}{1-\cos^2\theta - \sin\theta\cos\theta i} = \frac{1}{\sin^2\theta - \sin\theta\cos\theta i}$$

$$= \frac{1}{\sin\theta(\sin\theta - \cos\theta i)}$$

$$= \frac{1}{\sin\theta} \times \frac{\sin\theta + \cos\theta i}{\sin\theta - \cos\theta i}$$

$$= \frac{1}{\sin\theta} \times \frac{(\sin\theta + \cos\theta i)(\sin\theta - \cos\theta i)}{\sin\theta - \cos\theta i}$$



$$= \frac{\sin\theta + \cos\theta i}{\sin\theta} = 1 + i\cot\theta$$

(ii) $2\arg(z+1) = \arg(z)$

$$z+1 = \cos\theta + 1 + i\sin\theta$$

$$\arg(z+1) = \tan^{-1}\left(\frac{y}{x}\right) = \tan^{-1}\left(\frac{\sin\theta}{\cos\theta+1}\right)$$

Consider the diagram following:



$\triangle ABC$ is isosceles, and $\angle ABC = 180^\circ - \theta$

$$\therefore \angle BAC = \frac{1}{2}\theta$$

$$\therefore \angle CAE = \frac{1}{2}\theta$$

$$\therefore \arg(z+1) = \frac{1}{2}\theta = \frac{1}{2}\arg(z)$$

$$\therefore 2\arg(z+1) = \arg(z)$$


(c) The point $P(2p, p)$ is some point on the line $y = \frac{1}{2}x$ where $p \geq 0$.

- (i) Consider the locus of points that are the **same distance** from P as they are from the line $y = -2x$.

Explain why this locus is a parabola AND clearly describe its key features.

Note: you do **not** need to find the equation of the parabola.

~~$y = -2x$ is perpendicular to $y = \frac{1}{2}x$, hence, the~~
The locus represents ~~a~~ points equidistant from a line and a point, which defines a parabola




The vertex of the parabola is $(p, \frac{1}{2}p)$ and the directrix is $y = -2x$

- (ii) Consider another point $Q(q, -2q)$ on the line $y = -2x$ where $q \leq 0$.

Let R be the region enclosed by the locus of points that are **three times as far** from P as they are from the **point** Q .

Now suppose the points P and Q are moving along their respective lines.

If p is increasing at a rate of 3 cm s^{-1} , and q is decreasing at a rate of 2 cm s^{-1} , at what rate is the area of the region R increasing when P is $(6, 3)$ and Q is $(-4, 8)$?



$$\frac{dp}{dt} = 3, \frac{dq}{dt} = -2$$

~~R is a parabola with foci $(2p, p)$ and $(q, -2q)$~~

~~R should be a hyperbola - area is infinite as a hyperbola is unclosed~~

$$(x - q)^2 + (y + 2q)^2 = 3(x - 2p)^2 + (y - p)^2$$

$$x^2 - 2qx + q^2 + y^2 + 4qy + 4q^2$$

$$x^2 - 2qx + q^2 + y^2 + 4qy + 4q^2 = 3x^2 - 12px + 12p^2 + 3y^2 - 6py + 3p^2$$

$$2x^2 - (12p + 2q)x + 12p^2 - q^2 + 2y^2 - (6p + 4q)y + 3p^2 - 4q^2 = 0$$

Extra space if required.
Write the question number(s) if applicable.

QUESTION
NUMBER

1. a) i) When $x = \frac{1}{4}$,

$$\frac{d^2y}{dx^2} = -\frac{512}{225} < 0, \therefore \left(\frac{1}{4}, -\frac{2}{5}\right) \text{ is a local maxima}$$

when $x = -2$,

$$\frac{d^2y}{dx^2} = \frac{1}{18} > 0, \therefore (-2, \frac{1}{2}) \text{ is a local minima}$$

$$3. c) ii) \sin 2\theta = \frac{\frac{2abc}{b^2-a^2}}{c + \frac{2ca^2}{b^2-a^2}} = \frac{2abc}{b^2-a^2+2ca^2} = \frac{2ab}{c^2} = \frac{2abc}{c^3}$$

$$b^2c - a^2c + 2ca^2 = c^3$$

$$b^2 - a^2 + 2a^2 = c^2$$

$$a^2 + b^2 = c^2$$

$$3. b) i) \theta = 0; x = 5, y = 0, r = 5$$

$$\theta = \frac{\pi}{3}; x = \frac{3}{2}, y = \frac{3\sqrt{3}}{2}, r = 3$$

$\therefore$ Closest point in 1st quadrant is at $\left(\frac{3}{2}, \frac{3\sqrt{3}}{2}\right)$

$$3. b) ii) \frac{dx}{d\theta} = -4\sin\theta - 4\sin 4\theta = -4(\sin\theta + \sin 4\theta)$$

$$\frac{dy}{d\theta} = 4\cos\theta + 4\cos 4\theta = 4(\cos\theta + \cos 4\theta)$$

$$L = \int_0^{2\pi} \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta$$

$$= \int_0^{2\pi} 4 \sqrt{\sin^2\theta + 2\sin\theta\sin 4\theta + \sin^2 4\theta + \cos^2\theta + 2\cos\theta\cos 4\theta + \cos^2 4\theta} d\theta$$

$$= 4 \int_0^{2\pi} \sqrt{2 + 2\sin\theta\sin 4\theta + 2\cos\theta\cos 4\theta} d\theta$$

Extra space if required.
Write the question number(s) if applicable.

QUESTION
NUMBER

$$= 4 \int_0^{2\pi} \sqrt{2 + \cos \theta} d\theta$$

$$= 4 \int_0^{2\pi} \sqrt{2 + \cancel{2}\cos(-3\theta) - \cancel{2}\cos 5\theta + \cancel{2}\cos 5\theta + \cancel{2}\cos(-3\theta)} d\theta$$

$$= 4 \int_0^{2\pi} \sqrt{\cancel{4}2 + 2\cos(-3\theta)} d\theta$$

$$= 4 \int_0^{2\pi} \sqrt{2 + 4\cos^2\left(\frac{3}{2}\theta\right) - 2} d\theta$$

$$= 4 \int_0^{2\pi} 2\cos\left(-\frac{3}{2}\theta\right) d\theta$$

$$\cancel{2\pi} = \cancel{4} \int_0^{\frac{5\pi}{3}}$$

$$= 4 \times 3 \times \int_{\pi}^{\frac{5\pi}{3}} 2\cos\left(-\frac{3}{2}\theta\right) d\theta$$

$$= 24 \left[\frac{4}{3} \sin\left(\frac{3}{2}\theta\right) \right]_{\pi}^{\frac{5\pi}{3}}$$

$$= 64$$

2.0) ii) This is the minimum because it ensures that half of the absolute value components are inverted and of $f_n(x) = |x-n| = -x+n$ and half of $f_n(x) = |x-n| = x-n$, ensuring a minimum

Outstanding Scholarship

Subject: Calculus

Standard: 93202

Total score: 30

Q	Score	Marker commentary
1	7	The candidate exhibited elegance in presenting their work with logical clarity, demonstrating both a keen understanding and a commitment to coherence. They successfully recognised the need to calculate the differences between the heights of the stair steps I Q1(c), assuming the cubic curve reaches its turning point at the top of the hill.
2	8	The candidate demonstrated a comprehensive grasp of absolute value functions by adeptly expressing the summation of the series using symmetry in Q2(c). They successfully solved problems even when provided with minimal descriptions.
3	8	They demonstrated consistency in tackling proof-style questions in Q3(c) by following through each step until the completion of the task, displaying determination and perseverance.
4	7	The candidate showed the ability to transform a complex number problem into a simple geometry problem in Q4(b)(ii), showcasing strong analytical skills and a deep understanding of the interplay between two fields of mathematics.