

No part of the candidate's evidence in this exemplar material may be presented in an external assessment for the purpose of gaining an NZQA qualification or award.

SUPERVISOR'S USE ONLY

S

93104



Draw a cross through the box (☒) if you have NOT written in this booklet

☐

+

OUTSTANDING SCHOLARSHIP EXEMPLAR



Mana Tohu Mātauranga o Aotearoa
New Zealand Qualifications Authority

Scholarship 2023 Earth and Space Science

Time allowed: Three hours
Total score: 24

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

You should answer ALL the questions in this booklet.

Pull out Resource Booklet 93104R from the centre of this booklet.

Show ALL working.

If you need more room for any answer, use the extra space provided at the back of this booklet.

Check that this booklet has pages 2–16 in the correct order and that none of these pages is blank.

Do not write in any cross-hatched area (AREA NOT TO WRITE). This area may be cut off when the booklet is marked.

**YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE
END OF THE EXAMINATION.**

QUESTION ONE: PHYTOPLANKTON AND THE CARBON CYCLE

Discuss the significance of phytoplankton compared to plants in the global carbon budget.

In your answer, include how it could be affected by climate change, and how it could be enhanced using human intervention.

Both phytoplankton and plants play an important role in the Earth's carbon cycle.

~~Phytoplankton~~

Phytoplankton are what is part of the biological pump (which lies in addition to the physical pump), ~~and~~ with atmospheric CO_2 removed from the atmosphere ^{by dissolving in the water} where it is used in photosynthesis by phytoplankton. This carbon used in photosynthesis is used to produce organic forms of carbon such as glucose and other sugars, useful for respiration. ~~The oxygen can also~~ and the synthesis of other organic carbon molecules. Upon death, their bodies can also fall to the bottom of the ocean, the carbon being stored in limestone which after further compression and heat, after millions of years form coal, oil, gas, etc. These sediments are also able to be ~~sub~~ subducted ~~and as a way~~ with CO_2 again released into the atmosphere as part of the tectonic plates ~~with~~ and the formation eruption of volcanos, thereby playing a part in the rock cycle. After forming sediment and rock, they can eventually become part of the crust that subducts at a tectonic plate boundary to form subduction volcanoes. In this way, the death of phytoplankton leading to the

formation of ~~sediment~~ like limestone can act as a very long-term store of carbon. Similarly the carbon-rich bodies of phytoplankton ^{or other marine} can ^{life, like snakes which feed on other phytoplankton} sink to the deep ocean with their carbon sequestered. As there is very little mixing between the different layers of the ocean, this can also act as a long-term store of carbon, with this carbon ~~never even~~ ^{not} reaching the atmosphere for a long time as it slowly travels with the thermohaline current. This carbon will only ever reach the atmosphere ~~when~~ until travelling to sites of upwelling, thus being able to store carbon for long periods of time. This contrasts plants which are unable to contribute as significantly to phytoplankton in this long-term storage of carbon (though their dead bodies have the potential to form fossil fuels).

Both phytoplankton and plants will remove ~~atmosph~~ CO_2 from the atmosphere, and ~~are~~ ^{are} placed at the bottom of the food ~~as~~ ^{as} producers. They both are able to capture energy from the sun to allow photosynthesis to be performed. In the oceans, zooplankton eat phytoplankton which use energy stored in the organic forms of carbon produced by photosynthesis to perform photosynthesis releasing carbon dioxide:

$$\text{glucose} + \text{oxygen} \rightarrow \text{carbon dioxide} + \text{water} + \text{energy}$$

This CO_2 produced by respiration is then able

to return to the atmosphere. The same can be said for organisms that eat zooplankton and those further up the food web — essentially the entire ~~now~~ food web. ~~Deep~~ Dead phytoplankton whose bodies are not ^{stored} ~~absorbed~~ in deep ~~ocean~~ ^{ocean}, or the thermohaline currents or in limestone, along with all other ~~ocean~~ ~~decomposition~~ ~~on~~ dead marine organisms can further have their bodies decomposed by bacteria, also producing CO_2 that can return to the atmosphere. Plants, similar to phytoplankton also act as short-term stores of carbon in this way. ~~It~~ Organic material formed by photosynthesis is used by other living animals for respiration, the respiration of the plants themselves, or the respiration of bacteria feeding upon the dead ~~ocean~~ bodies of organic matter of organisms, all once again emitting CO_2 into the atmosphere again.

Coal, oil, gas, etc. are long-term carbon stores which ~~act as~~ ~~the~~ are formed from organic matter from millions of years ago which have undergone heat, compression, etc. As humans uncover these fossil fuels and burn them, they release carbon dioxide into the atmosphere — carbon that was previously stored. Increased carbon dioxide is ~~the~~ leading to climate change, with CO_2 acting as a greenhouse gas that is able to absorb

and reflected long-wave IR. This is leading to increased global temperatures. Increased CO_2 in the atmosphere may possibly also lead to more algal blooms, with large amounts of nutrients available and an excess of atmospheric CO_2 allowing large amounts of photosynthesis and further respiration which will lower O_2 contents in the water as it is used up in respiration. This, however, can block sunlight leading to the death of other marine producers ^{who can not photosynthesize} and cause the death of other marine wildlife with reduced O_2 levels for respiration.

Warming of the oceans will also lead Earth in climate change may also lead to the poles becoming significantly warmer, with the melting of sea ice, further meaning less short wave light is reflected from the sun as sea ice has a greater albedo than the darker ocean water, a helping a positive feedback loop. The melting of sea ice and increase in water temperature may all contribute to less downwelling of cold, saline water which could stop the deep thermohaline current. The true effects, however, are unknown, and this could possibly ~~also lead to~~ ^{have} the effect the deep ocean currents as a store of carbon. Warmer oceans may also lead to faster interactions in the marine ecosystem. A warmer ocean could mean ~~more photosynthesis~~ ^{the} faster breakdown of dead organic matter with a greater rate of photosynthesis, accelerating the release of CO_2 into

QUESTION TWO: SLOW-SLIP EVENTS

Explore the differences between earthquakes and slow-slip events, referring to the Eketāhuna and Kaikōura earthquakes.

In your answer, include how the information collected by geologists can help further their understanding of slow-slip events, and the difficulties with monitoring on land, and at sea.

In the North Island, as the denser, older oceanic crust of the Pacific Plate subducts under the less dense, newer, continental crust of the Australian plate, friction between the two plates causes the build-up of elastic potential energy ^{along the ~~overlying~~ ^{underlying} Australian plate} along faults. It is the sudden release of this elastic potential energy that results in the formation of an ~~big~~ earthquake with this energy ~~released~~ ^{released} all at once, whilst the gradual release is a slow-slip event.

Slow slip events occur when the Australian and Pacific plates move past each other slowly, releasing the elastic potential energy built up through friction over a longer period and larger area. This contrasts earthquakes which are more so the sudden release of energy ~~over a~~ in a short amount of time over a small area. Slow-slip events ^{also} occur at regular intervals, ~~which contrasts to earthquakes~~. During slow-slip events, the land moves very slowly as it slowly ~~releasing~~ releases the elastic potential energy built up from friction in the areas that have locked the tectonic plates together, and thus can take large amounts of time and be extremely slow moving as evidenced by the 2011

movement of the land. Because energy is released over a longer time period and much greater ones compared to Earthquakes, they are unlikely to cause much damage.

Yet ~~the~~ slow-slip events have also been linked to earthquakes. Earthquakes are likely to occur more in the locked plate interface zones with the plates locked by friction ~~preventing~~ ~~slow slipping~~ ~~preventing~~ slow-slip events and rather having large sudden fault movements. There are 3 types of faults: normal faults where crust is stretched, reverse where crust is ~~compressed~~ ~~compressed~~ compressed, and slip-strike faults which involve more lateral sideways movement of the faults.

A slip-strike ~~slow-slip events may result from~~ ~~Earthquakes as it causes~~ ~~disaster~~ Earthquakes such as the Eketahuna ~~earthquake~~ earthquake are thought to be the ~~result~~ ~~of~~ result of slip-strike slow-slip events such as the Kāpiti slow-slip. This ~~and~~ ~~Earthquake~~ ^{earthquake} lies on the boundary of the locked plate interface and slow slip zones, and may be caused by further stretching and build-up of elastic potential energy in the crust which ~~cause~~ ~~caused~~ ~~the~~ ~~fault~~ and thus resulted in the Eketahuna earthquake, ~~with~~ the plates snapping back into place releasing significant amounts of

energy. The 2016 Kaikōura earthquake has also been linked to slow-slip events that occurred after. For a similar reason, as the fault moves resulting in an earthquake, this could also cause a slow-slip event with the energy held in the plates, deformation along the fault causing a slow-slip event to release energy on this part of the crust. All of this motion and energy will build up on the overlying Australian plate as it rubs against the subducting Pacific Plate.

~~Slow Slip Earthquakes, the slow~~ Unlike earthquakes, as slow-slip events do not release large amounts of concentrated energy, they are unable to be able to be detected by seismometers with no noticeable shake in the ground. Therefore, they must be measured using GPS stations which will track the displacement of land. However, as the movement in land is ~~very~~ often extremely small, such as the 2cm displacement off the east coast near Pororua, these GPS stations are required to be extremely ~~difficult~~ precise and accurate, making taking measurements difficult. Similarly, it may be difficult to ~~do~~ However, it is shown that slow-slip events occur at regular intervals, like that

off the east coast near Pōrenghau with an event in 2006, 2011, 2016, and 2021 — an interval of every 5 years. Equipment can then be prepared to measure these slow-slip events in advance online with ~~Earthquakes~~ earthquakes which seemingly occur at more random intervals. To monitor land displacements accurately, numerous GPS stations will be needed and over a large area to measure the full extent of the slow-slip, also ~~over~~ with the slow slip zones comprising a large area proportion of New Zealand.


To ~~make~~ take recordings of slow-slip events at sea, pressure sensors are used to measure increases in height of the sea floor. This is as, when the Australian Plate slowly overrides the subducting Pacific Plate during slow-slip events, it will gradually uplift. This uplift means less water above the pressure sensor and so a lower pressure. Having to use pressure sensors, however, will come with many difficulties. It is likely very expensive to transport pressure sensors to the bottom of the sea floor, and changes in water pressure could be affected by numerous other factors like winds, or La Niña events for instance which would pile water on the east coast of NZ, raising the sea level and so pressure upon the sensor. ~~Diff~~ Differences in pressure would somehow need to be adjusted to measure only differences from the slow-slip event. Similarly, from pressure differences ~~also~~

QUESTION THREE: JUPITER'S ATMOSPHERIC CELLS

Analyse the differences between the Earth's atmospheric cells and the bands on Jupiter. Explain why these differences occur, and how these relate to the size, rotational speed, and composition of both planets.


In your answer, include a comparison of the depth of the cells between the two planets, and discuss the significance of the Giant Red Spot.

Whilst on Earth, there are only 6 cells, on Jupiter, there are 20 bands^{cells} which are similar to Earth's atmospheric cells. The reason for this is possibly due to the greater rotational speed of Jupiter which causes the Coriolis effect to have a larger impact. This means more deflection of winds to the right in the Northern hemisphere and more deflection to the left in the Southern hemisphere, with this greater deflection resulting in the need for more cells. Jupiter's bands are also much larger in size compared to Earth. This could be due to greater warming due to a larger surface area that means Jupiter receives more solar energy compared to Earth. On Earth, ~~and possibly Jupiter~~ the equator receives more solar radiation per unit area compared to the poles.



This causes an uneven heating effect that drives the polar and Hadley cells.

whilst the ferrel cells are not so much temperature driven. At the equator, warm, less dense air rises and falls at about 30° latitude North and South to form the Hadley cells. The other cells are also shown below.



The subtropical and polar jetstreams are also labelled which are areas of fast moving winds due to large temperature gradients between the cells. As Jupiter has more bands, as it spins faster, it will also have more of these jet streams. ^{that} are possibly faster as well, with a greater deflection east/west. ^{the jet streams are also faster.}

Jupiter's bands may also have a greater depth compared to Earth due to more solar radiation reaching it as well as as well as being able to produce more thermal radiation than it receives from the sun. ~~As a result~~ Unlike Earth, there is no solid rock that prevents larger convection currents being made up of mostly gas and liquid. Hence, this will allow much deeper convection currents to form with cool material sinking to the planet's centre, being warmed by the heat produced in the core as Jupiter cools from radioactive decay. ~~may~~ likely the heat left

over when Jupiter first formed. This causes the ~~gas material~~ now heated warm material which is less dense to rise, producing these large circulation cells, ^{back} compact to Earth in which, these circulation cells are driven only from solar energy alone. ~~As the material has reacted~~ ~~slowly on Jupiter, moving away from the equator~~ the cells may. Similar to Earth, Jupiter has Ferrell-like cells with circulation directions that are opposite to the temperature differences. As it is to As Jupiter has more bands it makes sense it would have more of the Ferrell-like cells compared to Earth, with one Ferrell-like cell between each temperature driven band. On both Earth and Jupiter vertical convective motion drives the bands/cells. ~~As the material~~ is heated, becoming less dense and so rises, until cooling, becoming more dense and falling until becoming heated again.

The Giant Red Spot is a storm on Jupiter on of crimson-colored clouds spinning anticyclonically at high speeds. It shows how Jupiter has far greater pressure gradients compared to Earth which can lead to such long-lasting and significant storms. Earth for ^{instance} ~~instance~~ has very short-lived and much weaker storms. It also proves how the bands on Jupiter and jets/wind movements are overall much faster.

and violent compared to Earth.

Overall, the atmospheric cells or bands on Jupiter are significantly larger with a greater depth and ~~fast than on Earth, meaning~~ are faster compared to Earth. ~~Convection~~ Vertical convection cells require fluids - liquids or gases that are free to move and make up the entirety of Jupiter, whilst only a small part of Earth which has a surface. This allows for deep vertical convection currents with material being significantly heated at the core of Jupiter, and cooling significantly as it reaches the surface and falls again. The strong temperature gradients as a result of this significant warming and cooling in addition to solar energy from the sun (though this is likely not as significant in driving the convection cells) contributes to these lat belts or jetstream equivalents on Earth ~~being~~ as a result of the massive pressure differences which drive these belts in a similar process to the polar or subtropical jet streams on Earth. The added 20 belts are ~~all~~ compared to Earth's 6 ~~atmospheric~~ cells are due to a more pronounced Coriolis effect with greater deflection ~~on~~ of air ~~to the right in the~~ to the right in the Northern hemisphere and left in the Southern hemisphere, thus requiring more atmospheric cells, and thereby in doing so leading to more ~~stronger~~ jet streams and intense Ferrel-like cells. In addition to showing how strong the temperature gradients are on Jupiter, the Great Red Spot may also demonstrate how strong the Coriolis force is on ~~the~~ Jupiter, producing such a

Back Page

Warmer water also absorbs less CO_2 than colder water, so oceans warming could lead to less CO_2 absorbed and less available for photosynthesis by phytoplankton leading to less by result.

Extra space if required.

Write the question number(s) if applicable.

QUESTION
NUMBER

- 1) the atmosphere. In terms of plants on land, climate change may have a similar effect. Warming may lead to a greater frequency of forest fires which would release large amounts of CO_2 into the atmosphere, in the burning of plants, with these plants also being now unable to absorb CO_2 from the atmosphere. Usually, the carbon cycle acts as the Earth's 'thermostat', with more CO_2 in the atmosphere leading to more rainfall (as warmer air can hold more water vapour) which removes CO_2 leading to less atmospheric CO_2 , so less rainfall, and so on. However, the burning of fossil fuels is placing large amounts of CO_2 that ~~is not~~ ^{is not} ~~being~~ ^{is} removed by this regulatory system is no longer in action. Human intervention to reduce the burning of CO_2 may not only ~~decrease atmospheric CO_2 concentrations~~ is also leading to burning of fossil fuels by adopting electric vehicles, etc. may ~~decrease~~ ^{reduce} CO_2 in the air and allow the regulatory system to once again act.



Increased atmospheric CO_2 is also leading to ocean acidification (or rather the oceans becoming less basic). Increased atmospheric CO_2 means more

Extra space if required.

Write the question number(s) if applicable.

QUESTION
NUMBER


CO₂ being absorbed to produce carbonic acid:



H⁺ produced from carbonic acid can react with CO₃²⁻ in the ocean, reducing the CO₃²⁻ in the ocean:



~~It~~ This makes it more difficult for carbonate shells to form with lower CO₃²⁻ concentrations and other carbonate minerals. It also reacts with CaCO₃ shells.



With less shells forming, less sediment is produced which can fall to the ocean floor, so the ocean acts as a worse long-term carbon store.

In this way, human intervention to prevent further burning of fossil fuels will reduce the increase in CO₂, and both reduce the warming of oceans and ocean acidification, reducing these negative effects and allowing better storage of carbon in oceans and ~~on land~~ ^{on land}. It would allow phytoplankton to better uptake and store carbon in the oceans long-term.

- 2) alone, it may be difficult to get accurate cm readings at the extent of ships. Likewise, with such a large area over which slow-ships can occur, numerous pressure sensors would all be required which could be expensive.

Extra space if required.
Write the question number(s) if applicable.

QUESTION
NUMBER

- 3) large storm of circular moving clouds in an anticyclonic direction, possibly similar to the pattern of gyres on Earth. As there are numerous jet streams, and the atmospheric cells massive compared to Earth, they move matter around Jupiter at significant speeds. Due to the atmospheric cells on Jupiter being much deeper compared to Earth, the jet streams are much ^{deeper} ~~deeper~~ as well - up to 3000 km deep as a result of these pressure gradients extending over a massive length.

93104

Subject	Earth and Space Science Scholarship Outstanding	Standard	93104	Total score	20
Q	Score	Annotation			
1	7	Candidate has explored the carbon cycle and related it to the context. Their answer is well structured and provides good coverage of human interventions and the effects of climate change on phytoplankton and carbon cycle. However, they have not compared the significance of plants with phytoplankton well enough to receive an 8 on this question.			
2	6	Candidate has analysed the differences between earthquakes and slow-slip events. Furthermore, they have extrapolated the resource material to explain how these events are linked. They completed with examining the issues around monitoring of slow-slip events and aimed to give some ideas regarding solutions. The answer was awarded a 6 because it was well structured. However, it was not awarded a 7 because the answer required more exploration of how these events are linked for example why did the Gisborne slow slip occur after the Kaikoura earthquake.			
3	7	Candidate has analysed differences between Earth and Jupiter's atmospheres as well as how they are affected by aspects of each of the planets. They were awarded a 7 because they did not fully explore the significance of the Giant Red Spot.			